

# NT Immunisation Schedule

## Children (up to 19 years)

May 2019

Age	Disease	Vaccine Brand	Dose No.	Reconstitute	Administration	Notes
<b>Birth</b>	Hepatitis B	H-B-Vax II or Engerix-B paediatric	1	×	IMI	Give within 7 days of birth
<b>6 weeks</b>	DTPa-HepB-IPV-Hib	Infanrix hexa	1	✓	IMI	ALL CHILDREN
	Pneumococcal	Prevenar 13	1	×	IMI	ALL CHILDREN
	Rotavirus	Rotarix	1	×	ORAL	Dose 1 must be given by 15 weeks of age
<b>4 months</b>	DTPa-HepB-IPV-Hib	Infanrix hexa	2	✓	IMI	ALL CHILDREN
	Pneumococcal	Prevenar 13	2	×	IMI	ALL CHILDREN
	Rotavirus	Rotarix	2	×	ORAL	Dose 2 must be given by 25 weeks of age
<b>6 months</b>	DTPa-HepB-IPV-Hib	Infanrix hexa	3	✓	IMI	ALL CHILDREN
	Pneumococcal	Prevenar 13	3	×	IMI	Aboriginal children, Medically at risk, Premature infants born under 28 weeks
<b>6 months to under 5 years</b>	Influenza	See Influenza on <i>Additional Notes</i> for vaccination	Annually	×	IMI	ALL CHILDREN 2 doses with 1 month interval when receiving the vaccine for the first time in their life between 6 months to under 9 years and annually thereafter.
<b>12 months</b>	Measles-mumps-rubella	Priorix or MMR II	1	✓	SC OR IMI	ALL CHILDREN
	Meningococcal ACWY	Nimenrix	1	✓	IMI	ALL CHILDREN
	Pneumococcal	Prevenar 13	3 or 4	×	IMI	ALL CHILDREN
	Hepatitis A	VAQTA	1	×	IMI	Aboriginal children
	Hepatitis B	H-B-Vax II or Engerix-B paediatric	4	×	IMI	Premature infants under 32 weeks or under 2000g at birth
<b>18 months</b>	Measles-mumps -rubella-varicella	Priorix-tetra or ProQuad	2	✓	SC OR IMI	ALL CHILDREN
	Haemophilis influenzae type b	Act-HIB	4	✓	IMI	ALL CHILDREN
	DTPa	Infanrix or Tripacel	4	×	IMI	ALL CHILDREN
	Hepatitis A	VAQTA	2	×	IMI	Aboriginal children
<b>4 years</b>	DTPa-IPV	Infanrix IPV or Quadracel	5	×	IMI	ALL CHILDREN
	Pneumococcal	Pneumovax 23	1	×	IMI	Medically at risk
<b>5 - 15 years</b>	Influenza	See Influenza on <i>Additional Notes</i> for vaccination	Annually	×	IMI	Aboriginal children 2 doses, 1 month apart if receiving the vaccine for the first time in their life and aged 6 months to under 9 years.
<b>12 years</b>	Human Papillomavirus	Gardasil 9	1	×	IMI	ALL CHILDREN
	Human Papillomavirus	Gardasil 9	2	×	IMI	ALL CHILDREN minimum five month interval between doses 1 and 2
	Human Papillomavirus	Gardasil 9	3	×	IMI	Medically at risk (3 dose course: 0, 2 and 6 months)
	Adult/adolescent dTpa	Boostrix or Adacel	1	×	IMI	ALL CHILDREN
<b>15 - 19 years</b>	Meningococcal ACWY	Nimenrix	2	✓	IMI	ALL CHILDREN if not previously given
	Pneumovax	Pneumovax 23	1	×	IMI	Aboriginal children