

Immunisation Schedule Queensland

April 2019

Children and adolescents (10 to 19 years) Always check the Australian Immunisation Register (AIR) prior to vaccinating

Any additional information about the vaccines included in this schedule should be obtained from the online Australian Immunisation Handbook (AIH) at www.beta.health.gov.au/health-topics/immunisation

Age	Disease	Vaccine brand	Important notes
School Immunisation Program – vaccinations not given in the school program should be caught up at the usual immunisation provider			
Year 7 students (or age equivalent)	Human papillomavirus (HPV)	Gardasil 9	<ul style="list-style-type: none"> • 2 doses given at 0 and 6 months • Immunocompromised adolescents require 3 doses given at 0, 2 and 6 months.
	Diphtheria-tetanus-pertussis	Boostrix	<ul style="list-style-type: none"> • 1 dose
Year 10 students (or age equivalent)	Meningococcal ACWY	Nimenrix <i>reconstitute</i>	<ul style="list-style-type: none"> • 1 dose
Additional vaccines for Aboriginal and Torres Strait Islander and medically at risk adolescents			
Aboriginal and Torres Strait Islanders (ALL age groups)	Influenza	As supplied	<ul style="list-style-type: none"> • 1 dose every year (Refer to the current Influenza Immunisation Program, Program Advice)
People with medical risk factors for invasive pneumococcal disease (IPD)	Pneumococcal	Pneumovax 23	Refer to the online version of the AIH for a list of medical risk factors and to determine who is eligible to be vaccinated

Age	Disease	Vaccine brand	Important notes		
Catch-up vaccines					
All 10 to 19 years (with partial or no documented history)	Diphtheria-tetanus-pertussis	Adacel OR Boostrix	<ul style="list-style-type: none"> • 3 doses at 0, 1 and 2 months • Minimum interval between each dose is 4 weeks 		
	Poliomyelitis	IPOL	<ul style="list-style-type: none"> • 3 doses at 0, 1 and 2 months • Minimum interval between each dose is 4 weeks 		
	Measles-mumps-rubella	Priorix OR MMRII <i>reconstitute</i>	<ul style="list-style-type: none"> • 2 doses required at 0 and 1 month • Minimum interval between each dose is 4 weeks 		
	Hepatitis B	H-B-VaxII OR Engerix B	10 to <11 years and 16 to 19 years <i>Use paediatric dose vaccine</i>	11 to 15 years <i>Use adult dose vaccine</i>	<ul style="list-style-type: none"> • 3 doses at 0, 1 and 6 months after dose • Minimum interval between dose 1 and dose 3 is 4 months
	Chickenpox	Varilrix OR Varivax <i>reconstitute</i>	<ul style="list-style-type: none"> • Only 1 dose required for adolescents <14 years • 2 doses required for individuals 14 years • Minimum interval between each dose is 4 weeks 		
Year 7 or age equivalent, up to 19 years	Human papillomavirus (HPV)	Gardasil 9	<15 years 2 doses	<i>Immunocompromised</i> 3 doses	15 to 19 years 3 doses
			<ul style="list-style-type: none"> • 0 (day of dose 1) • 6-12 months Note: If dose 1 is given before the 15th birthday only 2 doses are required	<ul style="list-style-type: none"> • 0 (day of dose 1) • 2 months • 6 months 	<ul style="list-style-type: none"> • 0 (day of dose 1) • 2 months • 6 months
All 15 to 19 years (with no previous dose)	Meningococcal ACWY	Nimenrix <i>reconstitute</i>	<ul style="list-style-type: none"> • 1 dose Note: If required NeisVacC can be provided for catch up on request		